

Prospective relationship between occupational physical activity and risk of ischaemic heart disease: are men and women differently affected?

Karen Allesøe ^{1,2*}, Mette Aadahl^{2,3}, Rikke Kart Jacobsen², Line Lund Kårhus², Ole Steen Mortensen^{1,4}, and Mette Korshøj¹

¹Department of Occupational and Social Medicine, Holbæk Hospital, part of Copenhagen University Hospital, Gl. Ringstedvej 4B, 4300 Holbæk, Denmark; ²Frederiksberg Hospital, Nordre Fasanvej 57, 2000 Frederiksberg, Denmark; ³Department of Clinical Medicine, University of Copenhagen, Blegdamsvej 3B, 2200 Copenhagen N, Denmark; and ⁴Section of Social Medicine, Department of Public Health, University of Copenhagen, Øster Farimagsgade 5, 1353 Copenhagen K, Denmark

Received 21 November 2022; revised 27 February 2023; editorial decision 1 March 2023; accepted 2 March 2023; online publish-ahead-of-print 8 March 2023

Aims

High occupational physical activity (OPA) seems to increase risk of cardiovascular diseases among men. However, findings are mixed, and it is not known if women are differently affected. Therefore, the aim of this study is to investigate the relationship between OPA and risk for ischaemic heart disease (IHD), and whether it differs across sex.

Methods and results

This prospective cohort study was based on 1399 women and 1706 men, aged 30–61 years, participating in the Danish Monica 1 study in 1982–84, actively employed, without prior IHD and answering an OPA question. The information on incidence of IHD, before and during the 34-years follow-up, was retrieved by individual linkage to the Danish National Patient Registry. Cox proportional hazards models were used to investigate the association between OPA and IHD. Compared to women with sedentary work, women in all other OPA categories had lower hazard ratio (HR) for IHD. Among men, the risk of IHD was 22% higher among those with light OPA, and 42% and 46% higher among those with moderate OPA with some lifting or strenuous work with heavy lifting, respectively, compared to men with sedentary OPA. Compared to women with sedentary work, HR for IHD was higher among men in all OPA categories. There was statistically significant interaction between OPA and sex.

Conclusion

Demanding or strenuous OPA seems to be a risk factor for IHD among men, whereas a higher level of OPA seems to protect women from IHD. This emphasizes the importance of taking sex differences into account in studies of health effects of OPA.

Lay summary

In the Danish Monica I study among 1399 women and 1706 men, we investigated whether high physical activity at work was associated with higher risk of ischaemic heart disease and whether this association was different among men and women.

The association between occupational physical activity and ischaemic heart disease was different among men and women.

High physical activity at work was associated with around 45% higher risk of ischaemic heart disease in men, but with around 65% lower risk in women.

The underlying mechanisms for this difference, e.g. differences in exposure and physiology, should be further investigated in future studies.

Keywords

Physical activity • Work • Heart diseases • Sex differences • Epidemiology

* Corresponding author. Fax: +45 38633977, Email: kalle@regionsjaelland.dk

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Introduction

Globally and in European countries, cardiovascular diseases (CVD) are among the leading causes of mortality.^{1,2} Furthermore, in 2017 in Europe, there were 108.7 million people living with CVD, hereof 34.9 million living with ischaemic heart disease (IHD).¹

Leisure time physical activity (LTPA) is well-known to promote health and to prevent many chronic diseases including CVD.³ However, the beneficial cardiovascular effect of physical activity may depend on the domain, i.e. whether physical activity is performed at work or during leisure time, and on the characteristics of performed activities.⁴ A recent review showed that whereas LTPA was protective, high occupational physical activity (OPA) did not protect from CVD mortality,⁵ and a recent study in a large population showed lower risk of major adverse cardiovascular events from LTPA but higher risk from high OPA.⁶ It is suggested that fundamental differences between LTPA and OPA including differences in intensity, amount of static vs. dynamic movements, duration, and time for recovery may be possible explanations of these seemingly opposing associations with health, the so-called physical activity health paradox.⁷

In studies analysing men and women separately, some studies among men showed detrimental effects of high OPA on cardiovascular health.^{8–12} Yet other studies did not find an association between OPA and heart disease, specifically, or CVD, including conditions affecting the heart and blood vessels, in men.^{13,14} Among women, a few studies showed¹⁵ or indicated^{9,16} that high OPA was associated with increased risk of heart disease, whereas other studies have indicated a protective effect or have shown no association.^{8,11,12,17} However, beneficial effects of OPA with respect to risk of heart disease and CVD have also been observed among both women^{18,19} and men,^{17–19} so overall the findings are mixed. Some studies have included both men and women and have analysed men and women separately, but very few have tested whether there is a sex difference.^{6,9,20} In a previous paper on gender and occupational health²¹ and in a recent paper on the future agenda for studies of the association between health and LTPA and OPA, respectively,²² one of the recommendations was to further explore the differences between men and women. Women have an average 7- to 10-year delay in the first manifestation of heart diseases compared to men,²³ and there are sex differences in basal metabolism, hormones, physical capacity, and muscle strength that may play a role in both the physical impact and perception of a given workload.^{24,25} Furthermore, there is extensive gender segregation in the labour market, especially in jobs with high physical demands,²⁶ and even though men and women have the same occupational titles, there are large differences in their physical working conditions.²⁷

The aim of the present study was to investigate the relationship between OPA and risk for IHD and possible differences in this relationship between men and women in the Danish Monica I (Monitoring Trends and Determinants of Cardiovascular Disease) study. Our hypothesis is that demanding or strenuous OPA is associated with a higher risk of IHD than lower levels of OPA, irrespective of sex.

Methods

Study population and exclusion criteria

The Danish Monica I study was conducted from 1982 to 1984. From the Danish Central Person Register, 4807 persons, hereof 2404 men and 2403 women from 11 municipalities in the western part of Copenhagen Region, born in 1922, 1932, 1942, or 1952, were drawn as a random selection and were invited to participate in the Danish Monica I health survey. Totally, 4807 were invited hereof 968 individuals aged 30, 980 aged 40, 965 aged 50, and 872 aged 60. The participation rate was 79%. After exclusion of those not actively employed at baseline, with missing information on OPA or with a hospital admission with IHD before baseline, the study

population consisted of 3105 respondents, hereof 1706 men and 1399 women (Figure 1).

Study design

This prospective cohort study was based on data from the Danish Monica 1 study with baseline at the date of health examination of each participant in 1982–84 and with register-based information on incident IHD during follow-up until 20 December 2016.

Endpoints

Information about incident cases of IHD was retrieved by individual linkage of a personal identification number assigned to all residents in Denmark to the Danish National Patient Registry²⁸ in the period from this registry was initiated in 1977 and until end of follow-up. Ischaemic heart disease was classified according to the WHO International Classification of Diseases (ICD 10, from 1994) and ICD 8 (from 1977 until the end of 1993), and IHD cases were defined as hospitalisation for myocardial infarction (410 in ICD-8 and I21–23 in ICD-10), other acute or chronic IHD (411–412 in ICD-8 and I24–25 in ICD-10), angina (413 in ICD-8 and I20 in ICD-10), or electrocardiographically diagnosed heart disease (414 in ICD-8).

Participants were censored as cases at their first event (first ever hospitalisation with IHD). Otherwise, they were censored when they died at or out of hospital, when they were classified as emigrated and could no longer be followed in the registers, and otherwise at the end of the follow-up. Information on date of death or classification as emigrated was retrieved from the Central Person Registry.

Assessment of occupational physical activity

High occupational physical activity was assessed by a single question, based on the Saltin and Grimby question.²⁹ Classification is shown in parenthesis:

To which of the following groups do you belong at your workplace—outside or at home:

- (1) Mainly sedentary: desk work, assemble small parts, and the like (sedentary)
- (2) Work involving some walking but no carrying heavy items: light industrial work, non-sedentary office work, inspection, kitchen work, housework, teaching, and the like (light)
- (3) Mainly walking, work involving climbing stairs, and some lifting: mail delivery, construction work, move heavy furniture, and the like (moderate, some lifting)
- (4) Physically demanding work with heavy lifting: excavation work, forestry, concrete work, and the like (strenuous, heavy lifting).

Assessment of covariates

Register based covariates: Sex (considered both a biological and a cultural/psychosocial factor) and age at the time of the health examination were retrieved from the Central Person Registry. Participants were included in the study according to four specific years of birth (30, 40, 50, and 60 years old at invitation to participate) but due to two years of inclusion time, the categorical age variable formed was: 30–32, 40–42, 50–52, and 60–62.


Covariates assessed at baseline investigations: At a health examination, clinical and biochemical data were collected, participants were interviewed, and a self-report questionnaire was filled out. For the assessment of medication use, the participants were asked to bring their medication used currently or during the last 12 months before baseline. The wording of the questions, measurement methods, and categorization of the covariates are found in section A in appendix.

Sociodemographic and socioeconomic factors (SES): Cohabitation status and vocational education.

Additional known risk factors for IHD: LTPA, self-reported fitness, smoking, alcohol intake per week, family history of heart disease, diabetes, body mass index (BMI), blood lipids: high density lipoprotein (HDL), serum cholesterol, and triglyceride.

Occupational factors: Work hours per week.

Variables for sensitivity analyses: Blood pressure, hypertension.



Statistical methods

In a current paper on perspectives for future research in the physical activity health paradox, it is recommended to investigate sex differences,²² hence it was decided *a priori* to do so, and to test for interaction between OPA and sex. This was done by inclusion of an interaction product term for OPA and sex (OPA × sex) and thereby allowing for the associations of OPA with the outcome to differ between sexes.

Population characteristics are presented by frequencies and percentages or mean and standard deviations (whenever appropriate) for the total population and for men and women separately, and according to level of OPA.

Due to the competing risks of IHD and death, risks of both are presented by cumulative incidence (CI) curves for men and women. The CI curves are compared by Gray's test for equality.³⁰

Cox proportional hazards (Cox PH) models were used to test for associations between OPA and IHD. In the models, follow-up time was the underlying time scale. In all Cox PH models, 95% confidence intervals (CI) were calculated. From available covariates, potential confounders for the association between OPA and IHD were selected *a priori* based on current knowledge from the literature about their association with CVD.^{31,32}

In the models without an interaction term, between OPA and sex, we analysed Model A: adjusted for age and Model B: adjusted for age and sex.

In a second model, the association between OPA and IHD was allowed to differ according to sex by including a term of interaction. The result from this model is presented both with a common reference group (women with sedentary OPA) and as the effect of OPA on risk of IHD within sex-strata. The following models were analysed: (i) Model 1: adjusted for age, (ii) Model 2: adjusted for age and LTPA, (iii) Model 3: Model 2 additionally adjusted for smoking, alcohol, BMI, self-reported fitness, diabetes, serum cholesterol, HDL, triglycerides, familial predisposition for heart disease, working hours, and civil status, and (iv) Model 4: Model 3 further adjusted for socioeconomic status (SES, vocational education).

Furthermore, the interaction between OPA and age, and OPA and LTPA was assessed.

The adequacy of the Cox proportional regression model was assessed by cumulative sums of martingale-based residuals.³³

As OPA is associated to SES,^{12,34} we made a sensitivity analysis of the age-adjusted association between OPA and IHD among those with the lowest level of vocational training: without or ≤ 1 years of training.

An increased systolic blood pressure (SBP) or hypertension may be in the causal pathway from high OPA to IHD, and therefore we performed a separate sensitivity analysis where Models 1 and 4 were further adjusted for SBP and made an analysis where Model 1 was stratified by hypertension.

Finally, as the follow-up time in the present study is long and participants may have changed exposure to OPA, e.g. due to retirement, the age-adjusted association between OPA and IHD was investigated in a model with 20 years of follow-up.

In all analyses, $P < 0.05$ was considered statistically significant. Statistical analyses were performed using the statistical package SAS version 9.4.

Results

The study population included 3105 respondents, 1706 men and 1399 women. Average follow-up time was 25.5 years (0.2–34.2 years) for men and 29 years (0.5–34.2 years) for women. The total number of person-years was 43 502.4 years for men and 40 529.3 years for women. During follow-up 358 men and 152 women were admitted to hospital with IHD. Of the remaining 1348 men and 1247 women, 594 men and 394 women died during follow-up, and 20 men and 12 women emigrated, or could otherwise no longer be followed in the registers. There were 734 men and 841 women who were followed until the end of follow-up.

Table 1 Characteristics (number and percentages or mean and standard deviation (SD)) among 1706 men and 1399 women participating in the Danish Monica 1 study

| | | All participants (n = 3105) | | | Women (n = 1399) | | | Men (n = 1706) | | |
|--|----------------------------------|--------------------------------|------|---------------|------------------|------|--------------|----------------|------|---------------|
| | | n | % | Mean (SD) | n | % | Mean (SD) | n | % | Mean (SD) |
| Sociodemographic and socioeconomic factors | | | | | | | | | | |
| Age | Mean age | | | 43.6 (10.3) | | | 42.8 (10.0) | | | 44.2 (10.5) |
| | 30–32 years | 895 | 28.8 | | 428 | 30.6 | | 467 | 27.4 | |
| | 40–42 years | 920 | 29.6 | | 430 | 30.7 | | 490 | 28.7 | |
| | 50–52 years | 840 | 27.1 | | 382 | 27.3 | | 458 | 26.9 | |
| | 60–62 years | 450 | 14.5 | | 159 | 11.3 | | 291 | 17.1 | |
| Civil status | Married/cohabitating | 2621 | 84.4 | | 1128 | 80.6 | | 1493 | 87.5 | |
| | Living alone | 484 | 15.6 | | 271 | 19.4 | | 213 | 12.5 | |
| Vocational training | Unskilled | 404 | 13.0 | | 245 | 17.5 | | 159 | 9.3 | |
| | Short (<2 years) | 2249 | 72.4 | | 992 | 70.9 | | 1257 | 73.7 | |
| | Medium theoretic (3–4 years) | 345 | 11.1 | | 139 | 9.9 | | 206 | 12.1 | |
| | Long higher education (>5 years) | 107 | 3.45 | | 23 | 1.6 | | 84 | 4.9 | |
| Occupational physical activity | | | | | | | | | | |
| OPA | Sedentary | 883 | 28.4 | | 458 | 30.7 | | 425 | 24.9 | |
| | Moderate | 1530 | 49.3 | | 739 | 52.8 | | 791 | 46.4 | |
| | Demanding | 526 | 16.9 | | 176 | 12.6 | | 350 | 20.5 | |
| | Strenuous | 166 | 5.4 | | 26 | 1.9 | | 140 | 8.2 | |
| Risk factors for IHD | | | | | | | | | | |
| LTPA | Moderate/vigorous | 663 | 21.4 | | 176 | 12.6 | | 487 | 28.6 | |
| | Light | 1577 | 50.8 | | 761 | 54.4 | | 816 | 47.8 | |
| | Sedentary | 865 | 27.9 | | 462 | 33.0 | | 403 | 23.6 | |
| Smoking | Current | 1831 | 59.0 | | 771 | 55.1 | | 1060 | 62.1 | |
| | Former | 563 | 18.1 | | 211 | 15.1 | | 352 | 20.6 | |
| | Never | 711 | 22.9 | | 417 | 29.8 | | 294 | 17.2 | |
| Alcohol intake ^a | Intake Units/week | | | 9.6 (11.9) | | | 5.11 (6.4) | | | 13.3 (13.9) |
| Self-rated fitness | Better than peers | 652 | 21.1 | | 218 | 15.6 | | 434 | 25.5 | |
| | Same as peers | 2184 | 70.5 | | 1055 | 75.5 | | 1129 | 66.4 | |
| | Worse than peers | 262 | 8.5 | | 124 | 8.9 | | 138 | 8.1 | |
| | Missing | 7 | | | | | | | | |
| BMI continuous ^a | kg/m ² | | | 24.6 (3.8) | | | 23.6 (4.0) | | | 25.3 (3.5) |
| Diabetes | No | 3046 | 98.1 | | 1380 | 98.6 | | 1666 | 97.6 | |
| | Yes | 59 | 1.9 | | 19 | 1.4 | | 40 | 2.4 | |
| AMI/other heart disease close relatives | No | 2208 | 72.1 | | 958 | 69.5 | | 1250 | 74.2 | |
| | Yes | 856 | 27.9 | | 421 | 30.5 | | 435 | 25.8 | |
| | Missing | 41 | | | | | | | | |
| Blood lipids ^a | Total cholesterol (mmol/L) | | | 6.02 (1.22) | | | 5.90 (1.2) | | | 6.12 (1.23) |
| | HDL cholesterol (mmol/L) | | | 1.49 (0.42) | | | 1.66 (0.42) | | | 1.35 (0.37) |
| | Plasma triglyceride (mmol/L) | | | 134.3 (111.6) | | | 109.5 (56.4) | | | 155.4 (138.1) |
| Blood pressure | Systolic blood pressure (mmHg) | 3105 | | 122.6 (16.1) | 1399 | | 119 (16.1) | 1706 | | 125.5 (15.4) |
| | Diastolic blood pressure (mmHg) | 3105 | | 76.8 (11.0) | 1399 | | 73.8 (10.6) | 1706 | | 79.3 (10.6) |
| Hypertension ^b | No | 2266 | 73 | | 1069 | 76.4 | | 1197 | 70.2 | |
| | Yes | 839 | 27 | | 330 | 23.6 | | 509 | 29.8 | |
| Occupational factors | | | | | | | | | | |
| Working time | Hours/week | 3097 | | 38.3 (9.8) | | | 32.9 (9.8) | | | 42.6 (7.4) |
| | Missing | 8 | | | | | | | | |

BMI, body mass index; AMI, acute myocardial infarct; OPA, occupational physical activity; LTPA, leisure time physical activity.

^aDue to anonymization, n is not shown, where the number of missing observations was under 5.

^bSBP \geq 140 or DBP \geq 90 or using antihypertensive medication or self-report of hypertension.

Table 2 Hazard ratio and 95% confidence interval for ischaemic heart disease according to level of occupational physical activity among 1706 men and 1399 women participating in the Danish Monica 1 study, 1982–84

| OPA | n/no. with IHD | Model A | | Model B | | Model C | |
|--------------------------|----------------|---------|--------------|---------|--------------|---------|--------------|
| | | HR | 95% CI | HR | 95% CI | HR | 95% CI |
| Sedentary | 883/133 | 1 | | 1 | | 1 | |
| Light | 1530/250 | 1.03 | 0.83 to 1.27 | 1.03 | 0.83 to 1.26 | 0.99 | 0.80 to 1.23 |
| Moderate, some lifting | 526/97 | 1.34 | 1.03 to 1.74 | 1.23 | 0.94 to 1.59 | 1.15 | 0.87 to 1.52 |
| Strenuous, heavy lifting | 166/30 | 1.49 | 1.00 to 2.22 | 1.18 | 0.79 to 1.75 | 1.14 | 0.75 to 1.74 |

A total of 358 cases among men and 152 among women of IHD during follow-up until December 2016.

Model A, adjusted for age; Model B, adjusted for age and sex; and Model C, adjusted for age, sex, leisure time physical activity, family history of heart disease, diabetes, body mass index (BMI), serum cholesterol, high density lipoprotein (HDL), triglycerides, smoking, alcohol consumption, self-reported fitness, working hours, civil status, and socioeconomic status (SES). HR, Hazard ratio; CI, confidence interval; IHD, ischaemic heart disease; OPA, occupational physical activity.

Table 1 presents the baseline characteristics of the study population among men and women. The mean age was 44.2 years among men, and 42.8 years among women.

Among women, those with sedentary OPA had the highest probability of experiencing IHD during most of the follow-up period. Women with light OPA had the lowest probability of IHD, except for strenuous OPA, where there were very few cases of IHD. Among men, those with strenuous OPA had the highest probability of IHD during most of the follow-up period. Men with sedentary OPA had the lowest probabilities of IHD during most of the follow-up period, especially in the last part of follow-up. There were no statistically significant differences according to Gray's test.

In Supplementary material online, Figure B in appendix is a figure of the CI curve of the probability of experiencing all-cause mortality—given no previous case of IHD—in each OPA category at any given time during follow-up. Among men, those with moderate OPA had the highest, and those with strenuous OPA had the lowest probability of all-cause mortality during most of the follow-up period. Among women, the curve shows that those with moderate OPA had the lowest probability of all-cause mortality during most of follow-up (Supplementary material online, Figure B in appendix).

Assessment of the adequacy of the Cox proportional regression model showed that the model assumption of proportional hazards was in accordance with data.

Table 2 presents the association between OPA and IHD among all participants. The age-adjusted HR for IHD was higher, the higher level of OPA, and was 1.49 (95% CI 1.00–2.22) for strenuous work compared to sedentary work in the age-adjusted analysis (Table 2). Further adjustment for sex especially attenuated the estimate for strenuous OPA, and adjustment for all covariates in Model C further attenuated the estimates (Table 2).

To investigate if the association between OPA and IHD differed by sex, an interaction term between OPA and sex was included, showing a statistically significant interaction between OPA and sex ($P = 0.048$ in Model 1 and $P = 0.032$ in Model 4).

There was no statistically significant interaction between OPA and age ($P = 0.672$), and OPA and LTPA ($P = 0.666$).

All models below included an interaction term between OPA and sex. Table 3 shows HR for IHD from different levels of OPA in strata of men and women with sedentary work as reference category for both sexes.

Among women, HR for IHD was lower in all other OPA categories, compared to sedentary work in the age-adjusted analysis, showing a 20–40% lower risk of IHD among those with light or moderate work. The lowest HR was seen among women with strenuous work, this category, however, was small (Model 1). Among men, the risk of IHD was 25% higher among those with light OPA, and 49% and 45%

higher among those with moderate or strenuous work, respectively, compared to men with sedentary OPA in the age-adjusted analysis. Adjusting for LTPA had only minor impact on the estimates (Model 2). Further adjustment for several known risk factors for IHD, working time, and civil status in Model 3 resulted in a slight attenuation of the magnitude of the associations among both men and women. Further adjustment for SES in Model 4 had only minor effects on the estimates (Table 3).

Table 4 shows results from the same model, but presented with a common reference group (women with sedentary OPA), to be able to see the combined effect of OPA and sex on risk of IHD. The estimates for women were the same as those in Table 3 in Models 1–4 as the reference group was the same. Among men, HR for IHD was higher in all OPA categories compared to women with sedentary OPA. In Models 1 and 2, men with sedentary work had nearly 50% higher risk of IHD, men with light OPA had around 90% higher risk, and men with moderate or strenuous OPA had more than two times higher risk of IHD, compared to women with sedentary work (Table 4). However, further adjustment in Model 3 especially attenuated the estimates among men. This was partly due to adjustment for HDL and to a lesser degree due to adjustment for BMI. The adjustment in Model 3 seemed to lower the sex difference in risk of IHD among sedentary men and women, and to a lesser extent the differences in risk of IHD between the other OPA categories among men and women, respectively. Further adjustment for SES in Model 4 only led to minor changes in the estimates. Further adjustment for SES in Model 4 only led to minor changes in the estimates. In Supplementary material online, Table C in appendix shows the same model with sedentary men as reference group (see Supplementary material online, Table C in appendix).

In a separate sensitivity analysis, Models 1 and 4 in Table 3 were further adjusted for SBP. This only resulted in minor changes of the estimates (results not shown). We also stratified for hypertension in a sensitivity analysis to see whether individuals with hypertension had higher risk of IHD from high OPA than normotensive individuals (see Supplementary material online, Table D in appendix). Among men, the largest difference between normotensives and hypertensives was among those with strenuous work, where HR for IHD compared to sedentary work was nearly twice as high among those with hypertension HR 1.97 (95% CI 0.99–3.92), but only 30% higher among normotensives HR 1.31 (95% CI 0.75–2.27). However, there were only 31 hypertensive men with strenuous work. Among women, a u-shaped association was indicated among hypertensive women with a lower risk of IHD among those with light OPA and higher risk of IHD among those with moderate OPA. The association with IHD among women with moderate OPA had the opposite direction according to hypertension status. However, among hypertensive women, there were only few

Table 3 Hazard ratio and 95% confidence interval for ischaemic heart disease according to level of occupational physical activity and sex in a model with an interaction term between occupational physical activity and sex

| Sex | OPA | No. of subjects/no. with IHD | Model 1 | | Model 2 | | Model 3 | | Model 4 | |
|--------------------------|--------------------------|------------------------------|-----------|--------------|-----------|--------------|-----------|--------------|-----------|--------------|
| | | | HR | 95% CI | HR | 95% CI | HR | 95% CI | HR | 95% CI |
| Women | Sedentary | 458 (57) | 1 | | 1 | | 1 | | 1 | |
| | Light | 739 (74) | 0.72 | 0.51 to 1.02 | 0.74 | 0.52 to 1.04 | 0.68 | 0.48 to 0.97 | 0.67 | 0.47 to 0.96 |
| | Moderate, some lifting | 176 (-) ^a | 0.83 | 0.50 to 1.38 | 0.86 | 0.52 to 1.44 | 0.77 | 0.46 to 1.29 | 0.77 | 0.46 to 1.30 |
| | Strenuous, heavy lifting | 26 (-) ^a | 0.35 | 0.05 to 2.55 | 0.37 | 0.05 to 2.68 | 0.30 | 0.04 to 2.17 | 0.30 | 0.04 to 2.17 |
| Men | Sedentary | 425 (76) | 1 | | 1 | | 1 | | 1 | |
| | Light | 791 (176) | 1.25 | 0.96 to 1.64 | 1.27 | 0.97 to 1.66 | 1.23 | 0.93 to 1.61 | 1.22 | 0.93 to 1.61 |
| | Moderate, some lifting | 350 (77) | 1.49 | 1.08 to 2.05 | 1.54 | 1.12 to 2.12 | 1.40 | 1.01 to 1.94 | 1.42 | 1.01 to 1.98 |
| Interaction sex x OPA | Strenuous, heavy lifting | 140 (29) | 1.45 | 0.94 to 2.22 | 1.48 | 0.97 to 2.28 | 1.44 | 0.92 to 2.26 | 1.46 | 0.93 to 2.29 |
| | | | P = 0.048 | | P = 0.052 | | P = 0.032 | | P = 0.032 | |

A total of 1706 men and 1399 women participating in the Danish Monica 1 study, 1982–84; 358 cases among men and 152 among women of IHD during follow-up until December 2016.

Model 1: adjusted for age; Model 2: adjusted for age and leisure time physical activity; Model 3: adjusted for age and leisure time physical activity, family history of heart disease, diabetes, body mass index (BMI), serum cholesterol, high density lipoprotein (HDL), triglycerides, smoking, alcohol consumption, self-reported fitness, working hours, civil status; and Model 4: Model 3 including socioeconomic status (SES).

HR, Hazard ratio; CI, confidence interval; IHD, ischaemic heart disease; OPA, occupational physical activity.

^aData protection regulations in Denmark require a minimum of 5 individuals in each group. Therefore we have collapsed number of cases in the moderate and strenuous categories among women. There were in total 21 cases of ischaemic heart disease (IHD) in the moderate and strenuous categories among women.

women with moderate and strenuous OPA and only few cases of IHD (see [Supplementary material online, Table D](#) in appendix). There was no statistically significant interaction between OPA and hypertension ($P = 0.732$).

An additional sensitivity analysis of the association between OPA and IHD among those with a low level of vocational training, without training, or ≤ 1 years of training, was performed. Among 623 women and 393 men with a low level of vocational training, the age-adjusted association between OPA and IHD was resembling, which is shown in [Table 3](#), at all levels of OPA with even stronger associations among men with moderate [HR 1.59 (95% CI 0.78–3.25)] or strenuous OPA [HR 1.66 (95% CI 0.76–3.61)].

Furthermore, we made a sensitivity analysis with 20 years of follow-up. This analysis showed the same pattern and was comparable to [Table 4](#) with a higher risk of IHD among men with higher levels of OPA and a lower risk among women with higher levels of OPA—compared to men and women with sedentary OPA, respectively.

Discussion

Main findings

We aimed to investigate the association between OPA and IHD and to elucidate whether men and women were differently affected.

The association between OPA and IHD differed by sex. Moderate and strenuous OPA was associated with around 45% higher risk of IHD among men and around 65% lower risk of IHD among women, in the fully adjusted model. This difference was observed both in the CI curves and in the results of the survival analyses. To directly compare the associations in men and women, we also presented the results with a common reference group. Compared to women with sedentary OPA, HR for IHD among men was higher in all OPA categories, and men with moderate or strenuous OPA had more than two times higher HR for IHD in the age-adjusted analysis.

Comparison with other studies

Two studies, including both sexes, are in accordance with the finding in the present study of an adverse association between high OPA and risk of IHD among men, and an indication of a protective effect of high OPA among women compared to those with sedentary work.^{8,12} Also, there are studies, among either men or women, in accordance with the present study. Several studies among men showed detrimental effects of high OPA on cardiovascular health.^{8–12} As well, beneficial effects of OPA, with respect to risk of heart disease and CVD, have been observed among women in previous studies.^{18,19}

In contrast, some previous studies among men showed no association between OPA and heart disease or CVD^{13,14}, or suggested a protective effect with respect to heart disease.^{17–19} Among women, a few studies showed,¹⁵ or indicated,^{9,16} that high OPA was associated with increased risk of heart disease, whereas other studies have shown no association.^{8,11,17} Though there are several studies in accordance with the findings of this study, overall, the findings are mixed.

A general problem with examining the association between OPA and IHD in women, in the working age, is that large populations are needed to have sufficient statistical power, as IHD is a rarer event among women than among men, in the working age.²³

Differences between men and women

The present study is one in very few demonstrating that the association between OPA and IHD differs by sex, shown by the statistically significant interaction between OPA and sex. In accordance, a study from 2013 also showed statistically significant interaction between OPA and sex.⁹ In contrast, two Danish studies did not find statistically

Table 4 Hazard ratio and 95% confidence interval for ischaemic heart disease according to level of occupational physical activity with a common reference group: women with sedentary occupational physical in a model with an interaction term between occupational physical activity and sex

| Sex | OPA | No. of subjects/ no. with IHD | Model 1 | | Model 2 | | Model 3 | | Model 4 | |
|-------|--------------------------|----------------------------------|---------|--------------|---------|--------------|---------|--------------|---------|--------------|
| | | | HR | 95% CI | HR | 95% CI | HR | 95% CI | HR | 95% CI |
| Women | Sedentary | 458 (57) | 1 | | 1 | | 1 | | 1 | |
| | Light | 739 (74) | 0.72 | 0.51 to 1.02 | 0.74 | 0.52 to 1.04 | 0.68 | 0.48 to 0.97 | 0.67 | 0.47 to 0.96 |
| | Moderate, some lifting | 176 (–) ^a | 0.83 | 0.50 to 1.38 | 0.86 | 0.52 to 1.44 | 0.77 | 0.46 to 1.29 | 0.77 | 0.46 to 1.30 |
| | Strenuous, heavy lifting | 26 (–) ^a | 0.35 | 0.05 to 2.55 | 0.37 | 0.05 to 2.68 | 0.30 | 0.04 to 2.17 | 0.30 | 0.04 to 2.17 |
| Men | Sedentary | 425 (76) | 1.48 | 1.05 to 2.08 | 1.57 | 1.11 to 2.21 | 0.99 | 0.68 to 1.47 | 0.99 | 0.67 to 1.46 |
| | Light | 791 (176) | 1.85 | 1.37 to 2.50 | 1.99 | 1.47 to 2.70 | 1.22 | 0.87 to 1.72 | 1.21 | 0.85 to 1.71 |
| | Moderate, some lifting | 350 (77) | 2.20 | 1.56 to 3.1 | 2.42 | 1.71 to 3.42 | 1.39 | 0.94 to 2.06 | 1.40 | 0.95 to 2.06 |
| | Strenuous, heavy lifting | 140 (29) | 2.14 | 1.37 to 3.35 | 2.32 | 1.48 to 3.64 | 1.44 | 0.88 to 2.35 | 1.44 | 0.88 to 2.36 |

A total of 1706 men and 1399 women participating in the Danish Monica 1 study, 1982–84; 358 cases among men and 152 among women of IHD during follow-up until December 2016. Model 1: adjusted for age; Model 2: adjusted for age and leisure time physical activity; Model 3: adjusted for age and leisure time physical activity, family history of heart disease, diabetes, body mass index (BMI), serum cholesterol, high density lipoprotein (HDL), triglycerides, smoking, alcohol consumption, self-reported fitness, working hours, civil status; and Model 4: Model 3 including socioeconomic status (SES).

HR, Hazard ratio; CI, confidence interval; IHD, ischaemic heart disease; OPA, occupational physical activity.

^aData protection regulations in Denmark require a minimum of five individuals in each group. Therefore we have collapsed number of cases in the moderate and strenuous categories among women. There were in total 21 cases of ischemic heart disease (IHD) in the moderate and strenuous categories among women.

significant interaction between OPA and sex.^{6,20} Otherwise, most studies that stratified by sex did not report tests for interaction between OPA and sex, or did not stratify by sex, but merely adjusted for sex. When aiming to establish whether the relationship between OPA and risk of IHD is in fact different in men and women, it is necessary to explore potential effect modification by sex.

Combining the results of examining the association between OPA and IHD, among both sexes, and the results in strata of sex demonstrates the importance of exploring sex differences in studies of health effects of OPA. In the analysis including both men and women, the opposite direction of the association between men and women is masked. One of the possible explanations of the sex difference, in the present study, may be that men and women have different levels of OPA, although their questionnaire answer placed them in the same OPA category. There is a high degree of gender segregation in the labour market, especially in jobs with high physical demands,²⁶ and a recent study showed that within sitting, standing, and walking occupations, the types of occupational groups were different for men and women.³⁵ Women in jobs with high physical demands are typically working in the health, social care, kitchen, or cleaning sectors.²⁶ This is also the case for lifting where a Danish study showed that women exposed to lifting were working in the health or cleaning sector, whereas men exposed to lifting held a variety of jobs.³⁶ Furthermore, even in the case where men and women have the same occupational titles, there are large differences in their physical working conditions.²⁷ In addition, the individual assessment of the physical demands at work, in a questionnaire, is partly subjective and the answers could be related to the physical capacity and health status of the respondent, and may also be relative to colleagues and others in the same profession.

Another possible explanation of the sex difference in the association between OPA and IHD could be physiological sex differences. Men and women are different with respect to e.g. muscle mass, physical capacity, and sex hormones, and these differences may play a role in both the physical impact and perception of a given workload.^{24,25} Furthermore, men and women at the same age differ in their vulnerability to heart disease due to a 7- to 10-years-later first manifestation of heart diseases in women.²³ Among women, menopause, and the

following changes in the level of sex hormones and the physiological consequences of this, may affect the association between OPA and IHD. It has been shown that the level of HDL is affected by menopause.³⁷ In the present study, adjustment for HDL affected the association between OPA and IHD in the analysis where women with sedentary OPA were the common reference group.

A preliminary analysis of the association between OPA and IHD stratified by menopause/use of hormone-replacement-therapy (HRT) and adjusted for age indicated that among menopausal women, using HRT, a higher level of OPA was associated with a higher risk of IHD, which was not the case among pre-menopausal or menopausal women not using HRT (results not shown). This emphasizes the importance of taking these factors into account in studies including menopausal women, and this finding should be further investigated in studies including more women.

Also, a sensitivity analysis of the association between OPA and IHD stratified for hypertension showed a different pattern among men and women with a higher risk of IHD among hypertensive men with strenuous OPA, than among normotensive men, and a tendency of a u-shaped association among hypertensive women, also observed in a previous study.³⁸ However, some of the strata were rather small, and this finding should be further investigated in larger cohorts.

Strengths and weaknesses of the study

The Monica study is a prospective study with a high response rate, it includes a fairly large number of both men and women, and it was therefore possible to explore sex differences. Information on IHD was obtained in a hospital-based registry, providing valid information on incident IHD²⁸ with almost complete coverage. Furthermore, this information is independent of information about the level of OPA.

It is a strength that the follow-up time is sufficiently long to obtain the number of cases of IHD necessary for statistical power. However, some workers may retire or change exposure during follow-up. If workers with high OPA to a higher degree, than those with light or sedentary OPA, change to jobs with less physical demands (the so-called healthy worker selection bias) or retire during follow-up, this could bias the results.




Figure 2 The cumulative incidence curves for hazard of both ischaemic heart disease at any given time during follow-up from baseline to end of follow-up among (A) women and (B) men.

However, a sensitivity analysis with 20 years follow-up time showed the same pattern as the main analyses. Two studies have shown that the level of OPA is rather stable over time.^{39,40} Future studies should investigate the impact of continuity and duration of OPA exposure.

The assessment of OPA is self-reported and based on the well-known Saltin and Grimby question,²⁹ and has been shown to be useful in population studies to divide the population in broad categories of OPA. However, it is a weakness of this question that it is not possible to separate walking and standing work. Standing work for many hours can impose strain on the cardiovascular system, and has been shown to

be associated with heart disease.³⁵ It would have been interesting to be able to separate these exposures.

The Monica studies were designed for studying risk factors for CVD, and therefore many relevant confounders related to risk of IHD are assessed. However, residual confounding from other relevant possible confounders, as psychosocial work factors and shift work, that were not assessed in the Monica study, cannot be ruled out.

The exposure and many of the covariates were assessed by self-report, which may have caused some degree of misclassification. This could dilute the size of the associations and lead to smaller effects.

High OPA is known to be associated with low SES^{12,34} and therefore it is important to take this into account in the analyses. Adjustment for SES only slightly changed the estimates in this study, and the sensitivity analysis of the association between OPA and IHD among those with short vocational training showed similar results as among all participants. This implies that SES is not the underlying explanation of the associations.

As an increased SBP may be in the causal pathway from high OPA to IHD, the impact of adjusting for SBP was investigated as a separate analysis. This adjustment had only minor impact on the estimates.

In conclusion, the association between OPA and IHD was different among men and women, and this study is one in very few to demonstrate this. Compared to sedentary OPA, demanding or strenuous OPA was associated with a higher risk of IHD among men. In contrast, a higher level of OPA seemed to be protective in relation to IHD compared to sedentary OPA among women. Future studies should investigate the underlying mechanisms for this difference, whether it is differences in exposure and/or physiological differences between the two sexes.

Supplementary material

Supplementary material is available at *European Journal of Preventive Cardiology*.

Author Contributions

All authors have made substantial contributions to the conception and design, acquisition of data, or analysis and interpretation of data; drafted the article or revised it critically for important intellectual content; given final approval of the version to be published; and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Acknowledgements

The authors thank the participants in the Monica I study, the staff members at the Center for Clinical Research and Prevention, for contributing to the data collection and health examinations and the researchers involved in the planning and execution of this population-based study.

Funding

Karen Allesøe and Mette Korshøj were supported by a grant from The Working Environment Foundation. Grant Number: 05-2019-03/20195100247

Conflict of interest: None declared.

Data availability

Data cannot be shared for ethical/privacy reasons.

References

1. Timmis A, Townsend N, Gale CP, Torbica A, Lettino M, Petersen SE, et al. European Society of Cardiology: cardiovascular disease statistics 2019. *Eur Heart J* 2020;**41**:12–85.
2. 2016 Causes of Death Collaborators. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;**390**:1151–1210.
3. Lollgen H, Bockenhoff A, Knapp G. Physical activity and all-cause mortality: an updated meta-analysis with different intensity categories. *Int J Sports Med* 2009;**30**:213–224.
4. Holtermann A, Hansen JV, Burr H, Søgaard K, Sjøgaard G. The health paradox of occupational and leisure-time physical activity. *Br J Sports Med* 2012;**46**:291–295.
5. Cillekens B, Huysmans MA, Holtermann A, van Mechelen W, Straker L, Krause N, et al. Physical activity at work may not be health enhancing. A systematic review with meta-analysis on the association between occupational physical activity and cardiovascular disease mortality covering 23 studies with 655 892 participants. *Scand J Work Environ Health* 2022;**48**:86–98.
6. Holtermann A, Schnohr P, Nordestgaard BG, Marott JL. The physical activity paradox in cardiovascular disease and all-cause mortality: the contemporary Copenhagen General Population Study with 104 046 adults. *Eur Heart J* 2021;**42**:1499–1511.
7. Holtermann A, Krause N, van der Beek AJ, Straker L. The physical activity paradox: six reasons why occupational physical activity (OPA) does not confer the cardiovascular health benefits that leisure time physical activity does. *Br J Sports Med*. 2017;**0**:1–2.
8. Holtermann A, Marott JL, Gyntelberg F, Søgaard K, Suadicani P, Mortensen OS, et al. Occupational and leisure time physical activity: risk of all-cause mortality and myocardial infarction in the Copenhagen City Heart Study. A prospective cohort study. *BMJ Open* 2012;**2**:e000556.
9. Hu GC, Chien KL, Hsieh SF, Chen CY, Tsai WH, Su TC. Occupational versus leisure-time physical activity in reducing cardiovascular risks and mortality among ethnic Chinese adults in Taiwan. *Asia Pac Public Health* 2014;**26**:604–613.
10. Krause N, Brand RJ, Arah OA, Kauhanen J. Occupational physical activity and 20-year incidence of acute myocardial infarction: results from the Kuopio Ischemic Heart Disease Risk Factor Study. *Scand J Work Environ Health* 2015;**41**:124–139.
11. Wanner M, Lohse T, Braun J, Cabaset S, Bopp M, Krause N, et al. Occupational physical activity and all-cause and cardiovascular disease mortality: results from two longitudinal studies in Switzerland. *Am J Ind Med* 2019;**62**:559–567.
12. Holtermann A, Burr H, Hansen JV, Krause N, Søgaard K, Mortensen OS. Occupational physical activity and mortality among Danish workers. *Int Arch Occup Environ Health* 2012;**85**:305–310.
13. Virkkunen H, Harma M, Kauppinen T, Tenkanen L. The triad of shift work, occupational noise, and physical workload and risk of coronary heart disease. *Occup Environ Med* 2006;**63**:378–386.
14. Sobolski J, Kornitzer M, De BG, Dramaix M, Abramowicz M, Degre S, et al. Protection against ischemic heart disease in the Belgian Physical Fitness Study: physical fitness rather than physical activity? *Am J Epidemiol* 1987;**125**:601–610.
15. Allesøe K, Holtermann A, Aadahl M, Thomsen JF, Hundrup YA, Søgaard K. High occupational physical activity and risk of ischaemic heart disease in women: the interplay with physical activity during leisure time. *Eur J Prev Cardiol* 2014;**22**:1601–1608.
16. Petersen CB, Eriksen L, Tolstrup JS, Søgaard K, Grønbaek M, Holtermann A. Occupational heavy lifting and risk of ischemic heart disease and all-cause mortality. *BMC Public Health* 2012;**12**:1070.
17. Graff-Iversen S, Selmer R, Sorensen M, Skurtveit S. Occupational physical activity, overweight, and mortality: a follow-up study of 47,405 Norwegian women and men. *Res Q Exerc Sport* 2007;**78**:151–161.
18. Hu G, Jousilahti P, Borodulin K, Barengo NC, Lakka TA, Nissinen A, et al. Occupational, commuting and leisure-time physical activity in relation to coronary heart disease among middle-aged Finnish men and women. *Atherosclerosis* 2007;**194**:490–497.
19. Barengo NC, Hu G, Lakka TA, Pekkarinen H, Nissinen A, Tuomilehto J. Low physical activity as a predictor for total and cardiovascular disease mortality in middle-aged men and women in Finland. *Eur Heart J* 2004;**25**:2204–2211.
20. Holtermann A, Marott JL, Gyntelberg F, Søgaard K, Mortensen OS, Prescott E, et al. Self-reported occupational physical activity and cardiorespiratory fitness: importance for cardiovascular disease and all-cause mortality. *Scand J Work Environ Health* 2016;**42**:291–298.
21. Messing K, Silverstein BA. Gender and occupational health. *Scand J Work Environ Health* 2009;**35**:81–83.
22. Coenen P, Huysmans MA, Holtermann A, Krause N, van Mechelen W, Straker LM, et al. Towards a better understanding of the 'physical activity paradox': the need for a research agenda. *Br J Sports Med* 2020;**54**:1055–1057.
23. Maas AH, Appelman YE. Gender differences in coronary heart disease. *Neth Heart J* 2010;**18**:598–602.
24. Ilmarinen J. Work and cardiovascular health: viewpoint of occupational physiology. *Ann Med* 1989;**21**:209–214.
25. Søgaard K, Sjøgaard G. Physiological bases of work assessment. In Wilson JR, Sharples S, eds. *Evaluation of Human Work*. 4. London: C R C Press LLC; 2015. p419–445.
26. Aittomaki A, Lahelma E, Roos E, Leino-Arjas P, Martikainen P. Gender differences in the association of age with physical workload and functioning. *Occup Environ Med* 2005;**62**:95–100.
27. Messing K, Tissot F, Saurel-Cubizolles MJ, Kaminski M, Bourguine M. Sex as a variable can be a surrogate for some working conditions: factors associated with sickness absence. *J Occup Environ Med* 1998;**40**:250–260.
28. Schmidt M, Schmidt SA, Sandegaard JL, Ehrenstein V, Pedersen L, Sorensen HT. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol* 2015;**7**:449–490.

29. Saltin B, Grimby G. Physiological analysis of middle-aged and old former athletes. Comparison with still active athletes of the same ages. *Circulation* 1968;**38**:1104–1115.
30. Gray RJ. A class of K-sample tests for comparing the cumulative incidence of a competing risk. *Ann Statist* 1988;**16**:1141–1154.
31. O'Flatherty M, Sans-Menendez S, Capewell S, Jørgensen T. Epidemiology of atherosclerotic cardiovascular disease: scope of the problem and its determinants. In: Gielen S, Backer DG, Piepoli MF, Wood D, eds. *The ESC textbook of preventive cardiology*. Oxford: Oxford University Press; 2015. p3–18.
32. Kivimaki M, Jokela M, Nyberg ST, Singh-Manoux A, Fransson EI, Alfredsson L, et al. Long working hours and risk of coronary heart disease and stroke: a systematic review and meta-analysis of published and unpublished data for 603,838 individuals. *Lancet* 2015; **386**:1739–1746.
33. Lin DYW, Ying Z. Checking the Cox model with cumulative sums of martingale-based residuals. *Biometrika* 1993;**80**:557–572.
34. Beenackers MA, Kamphuis CB, Giskes K, Brug J, Kunst AE, Burdorf A, et al. Socioeconomic inequalities in occupational, leisure-time, and transport related physical activity among European adults: a systematic review. *Int J BehavNutrPhysAct* 2012;**9**:116.
35. Smith P, Ma H, Glazier RH, Gilbert-Quimet M, Mustard C. The relationship between occupational standing and sitting and incident heart disease over a 12-year period in Ontario, Canada. *Am J Epidemiol* 2018;**187**:27–33.
36. Møller A, Reventlow S, Hansen AM, Andersen LL, Siersma V, Lund R, et al. Does a history of physical exposures at work affect hand-grip strength in midlife? A retrospective cohort study in Denmark. *Scand J Work Environ Health* 2013;**39**:599–608.
37. Karppinen JE, Tormakangas T, Kujala UM, Sipilä S, Laukkanen J, Aukee P, et al. Menopause modulates the circulating metabolome: evidence from a prospective cohort study. *Eur J Prev Cardiol* 2022;**29**:1448–1459.
38. Allesøe K, Søgaard K, Aadahl M, Boyle E, Holtermann A. Are hypertensive women at additional risk of ischaemic heart disease from physically demanding work? *Eur J Prev Cardiol* 2016;**23**:1054–1061.
39. Wang A, Arah OA, Kauhanen J, Krause N. Effects of leisure-time and occupational physical activities on 20-year incidence of acute myocardial infarction: mediation and interaction. *Scand J Work Environ Health* 2016;**42**:423–434.
40. Krause N, Brand RJ, Kaplan GA, Kauhanen J, Malla S, Tuomainen TP, et al. Occupational physical activity, energy expenditure and 11-year progression of carotid atherosclerosis. *Scand J Work Environ Health* 2007;**33**:405–424.